

Effect of *Hibiscus sabdariffa* extract on the body weight, complete blood count, liver function, and lipid profile in rabbits

Hadeel Tariq Saadoun Al-Shalash

Medical Laboratory Technology Department, Al-Noor University College, Nineveh, Iraq.

Abstract

The objective of the study conducted here was to investigate effect of *Hibiscus sabdariffa* extract on the body weight, complete blood count (CBC), liver function, and lipid profile in rabbits. A total of 20 local male rabbits were randomly divided into four experimental groups, including five rabbits in each group: (i) control; (ii) starvation; (iii) *H. sabdariffa* extract; and (iv) *H. sabdariffa* extract and starvation. Blood samples were collected directly from the heart at the end of the experiment (day 28) and used for complete blood count and estimation of cholesterol, triglycerides, ALT, and AST. Aorta and liver sections were histopathologically examined. Results indicated that the Body weight at the end of the study was significantly decreased in starvation, extract, and starvation and extract groups, compared to the beginning of the study for each group. There was no difference in the body weight between groups at the end of the study. Rabbits in the extract group showed significant increase in lymphocytes, RBC, hemoglobin, and platelets, compared to those in control and in starvation groups. ALT activity significantly increased in starvation, extract, and starvation and extract groups, compared to those in control group. AST activity significantly increased in both extract and starvation and extract group, compared to those in control and starvation groups. Cholesterol and triglycerides decreased in starvation, extract, and starvation and extract groups, compared to those in control group. Histopathological sections indicated simple changes in aorta layers in starvation group rabbits including few number foam cells in tunica media. Liver sections from starvation group rabbits showed coagulative necrosis of hepatocytes, hyperplasia of bile duct endothelia, dilation and congestion of central veins. In extract group there was dilation of central veins, vacuolar degeneration of the hepatocytes, infiltration of inflammatory cells and hepatic artery congestion. Similar changes were revealed in liver sections from starvation and extract group. In conclusion, aqueous *H. sabdariffa* can decrease the body weight as alternative to the starvation, decrease blood cholesterol and triglycerides, and increase the immunity; however, daily consumption can have impact on the liver.

Keyword: *Hibiscus sabdariffa*, starvation, rabbits, body weight, lipid, liver

INTRODUCTION

Hibiscus is a large genus plant from the family "Malvaceae" includes 200 to 220 species primarily cultivated for the use of the flowers, leaves, and seeds, and known as rosella or rosemallow (Hussein et al., 2010). Although hibiscus is a native to tropical Africa, it can also be found in Indian and Southeast Asia including Thailand and Malaysia (Mahadevan et al., 2009). Fresh or dried calyces are used for preparation of herbal drinks and various hot or cold beverages (Bako et al., 2010; Ismail and Nazri, 2008). In Egypt, Sudan and Nigeria, fresh calyces are used in preparing a drink known as Caredey tea, Karkade, or Zoborodo, respectively (Gibbor and Pain, 1985; Kochhar, 1986).

The nutritional composition of fresh calyces *Hibiscus sabdariffa* can vary depending on genetic, environmental, ecological, and harvest conditions of the plant (Da-Costa-Rocha et al., 2014). The leaf contains protein, fat, carbohydrate, fiber, ash, calcium, phosphor, iron, thiamine, β -carotene, riboflavin, niacin and ascorbic acid (Duke and Atchley, 1984). The calyces contain the flavonoids, alkaloids, β -sitosterol, anthocyanin, citric acid wax, pectin, crude protein and different minerals (Hirunpanich et al., 2006; Singh et al., 2017).

Hibiscus is considered safe medical plant (Abbas et al., 2011). Aqueous extract of *H. sabdariffa* petals exhibited antihypertensive and cardio-protective effects in rats (Odigie et al., 2003). Consumption of three servings of *H. sabdariffa* tea daily lowered the blood pressure in pre-and mid hypertensive adults (McKay et al., 2010). In addition, *Hibiscus sabdariffa* can have the antihypertensive effect via three major mechanisms: diuretic (Jiménez-Ferrer et al., 2012), vasodilator (Sarr et al., 2009), and angiotensin converting enzyme inhibitor (Ojeda et al., 2010). However, *H. sabdariffa* extract can reduce high blood pressure on long term as antioxidant (Chen et al., 2004), which considered a cardio-protective effect. In addition, hibiscus buds can remove burning sensations and relieve pain (Alka et al., 2009). The analgesic effect of the flower or leaves extracts is thought to be via inhibiting prostaglandin synthesis; however, the methanolic extracts endowed with analgesic effect mediated through central inhibitory mechanism (Zahid et al., 2014). Furthermore, *H. sabdariffa* can manage different medical conditions including cancer, inflammatory disease, and different cardiovascular problems (De-Casta-Rochal et al., 2014), as well as antibiotics resistance (Voon et al., 2012).

People are excessively use herbal plants, including hibiscus tea, particularly in some Arabic regions. However, effect of hibiscus on some body function is not well studied. The objective of this study was to investigate effect of *H. sabdariffa* extract with starvation on the body weight, complete blood count (CBC), liver function, and lipid profile in rabbits.

MATERIALS AND METHODS

Hibiscus sabdariffa extract preparation

Dry calyxes of *H. sabdariffa* were obtained from local market in Mosul, Iraq, and identified at the Herbarium of Botany Department, University of Mosul. A 10% aqueous extract was prepared as the following: 10 gram of dry *H. sabdariffa* calyx was added to 100 ml of boiled distilled water. The mixture was kept for 15 minutes at room temperature, and then filtered using Whatman filter paper, and stored at 4 °C.

Study Animals

A total of 20 local male rabbits 9 to 10 months old were used in this study. The rabbits were kept in an appropriate environment (temperature 28 °C, and 12 hrs dark/light). The rabbits were fed with a standard diet prepared locally and included 47% wheat bran, 38% crushed local barely, 10% soybeans, 2% protein concentrate, 1% limestone powder, 1.5% NaCl, and 0.5% minerals and vitamins mixture.

Experimental groups

The rabbits were randomly divided into four experimental groups, including five rabbits in each group: (i) control group (CG), where rabbits fed 3 times per day (8 am, noon, 6 pm) and received distilled water once daily at 6 pm (1ml water / 1kg BW); (ii) starvation group (SG) for 16 hrs; where rabbits were fed and received distilled water once daily at 6 pm (1ml water / 1kg BW); (iii) *H. sabdariffa* extract group (HG), where rabbits fed 3 times per day (at 8 am, noon, 6 pm) and received 10% aqueous *H. sabdariffa* extract once daily at 6 pm (1ml extract/ 1kg BW); and (iv) *H. sabdariffa* extract and starvation group (HSG), where rabbits were fed 1 time per day (6 pm) and received 10% aqueous *H. sabdariffa* extract (1ml extract/ 1kg BW). Water was open to all rabbits all the time. Rabbits in all groups were initially weighted and allowed one week of acclimatization before starting the experiment. In each groups, the rabbits received the treatment (distilled water or the extract) once daily at 6 pm for four weeks. At the end of the experiment, five ml of blood were collected from the heart directly, and then all rabbits were slaughtered by cervical dislocation. Liver and aorta (from the arch to the iliac bifurcation) were harvested.

Blood analysis

Blood EDTA samples were tested for complete blood count (CBC) including total red blood cells count, differential leukocytes count, hemoglobin, and platelets count using hematology analyzer (Siemens Healthcare Diagnostics, Eschborn, Germany). Blood serum samples were analyzed via cobas c111 analyzer (Roche DiaLog, New Zealand) to estimate cholesterol, triglycerides using commercial kits (Biocon, Bengaluru, India), and ALT and AST using commercial kits (Biolabo, Maizy, France)

Histopathological sections

Liver and aorta sections were prepared according to Drury et al (1985) including the following steps: washing, dehydration, clearing, infiltration and embedding, and trimming and sectioning. Finally, the sections were stained using hematoxylin and eosin (Luna, 1968).

Statistical Analysis

Differences between study groups were examined using one-way analysis of variance (ANOVA), and the difference between body weight at day 0 and 28 in each group was examined using t-test (Moore et al., 2009). The data were presented as Mean \pm Standard Deviation. The difference with value of $P \leq 0.05$ was considered statistically significant. Statistical analysis performed using STATA 13.0 (StataCorp, College Station, TX).

RESULTS AND DISCUSSION

Body Weight

Means of the body weight between the groups were not different at the beginning of the study ($P = 0.19$). Body weight at the end of the study (day 28) was significantly decreased ($P \leq 0.05$) in SG, HG, and HSG, compared to the beginning of the study (day 0) for each group. However, there was no difference in the body weight between groups at the end of the study ($P = 0.27$) (Fig 1).

In starvation, loss of body weight occurs due to loss of adipose tissue subcutaneous and around different internal organs in addition to muscle atrophy (Madea, 2005). In our study, treatment with aqueous *H. sabdariffa* extract decreased the body weight as same as for that for starvation. However, effect of *H. sabdariffa* extract on decreasing adipose tissue has not been examined, yet. A consumption of 500 mg per day of LC-HS for two months in significantly reduced body weight, abdominal circumference, and percentage of body fat overweight women and in obese volunteers consuming the polyphenolic extract (Herranz-Lopez et al., 2019; Al-Snafi, 2018). One explanation for *H. sabdariffa* extract decreased body weight is through reduce of dietary carbohydrate absorption; as the plant contains enzyme inhibitor blocks production of amylase, an important enzyme for carbohydrate break down (Singh et al., 2017). An additional possible explanation for the anti-obesity effect for *H. sabdariffa* is through the effect of the extract on leptin and ghrelin, as the extract increased glucagon-like peptide-1 and decreased ghrelin hormones (Herranz-Lopez et al., 2019).




Fig 1. Differences in body weight (kg) between study groups; control (CG); starvation (SG); *H. sabdariffa* extract (HG); and *H. sabdariffa* extract and starvation (HSG) groups, compared within groups and between day 0 (beginning of the study) and day 28 (end of the study). Different letters (i.e., a, b) refer to statistical difference ($P \leq 0.01$), star (*) means no statistical difference between the groups.

Hematology

Differences in hematological parameters between groups of the study were summarized in Table 1. Rabbits in HG group showed significant increase in RBC, hemoglobin, lymphocytes, and platelets, compared to CG and SG groups. Rabbits in HSG group did not differ from HG group, except for platelets. Hematological parameters in CG and SG groups were not different, except for platelets. Monocytes were not different between all groups.

One explanation for increase of RBC in HG is that *H. sabdariffa* extract might stimulate release of erythropoietin in the kidney, the humeral regulator of RBC production (Elsner et al., 2004). In addition, Kaur and Kapoor, (2005) showed that anthocyanins, flavonoids exist in *H. sabdariffa*, can induce renal secretion of erythropoietin. Another possible explanation is that bone marrow and lymphoid organs are stimulated by different compounds in *H. sabdariffa* calyx such as alkaloids, flavonoids, polyphenolics, ascorbic acid and other vitamins, and thus different blood cells are increased (Mungole and Chaturvedi, 2011).

Table 1. Differences in hematological parameters between study groups, control (CG); starvation (SG); *H. sabdariffa* extract (HG); and *H. sabdariffa* extract and starvation (HSG) groups, at the end of the study. Different letters (i.e., a, b, c, d) for each parameter refer to statistical difference ($P \leq 0.05$).

Hematological parameter	Mean \pm Standard deviation			
	CG	SG	EG	ESG
Red blood cells (cell $\times 10^6\mu\text{L}$)	4.29 \pm 0.57 ac	3.75 \pm 0.46 a	5.37 \pm 0.39 b	4.96 \pm 0.46 cb
Hemoglobin (g/dL)	7.2 \pm 0.22 a	7.08 \pm 0.73 a	8.5 \pm 0.58 b	8.36 \pm 0.38 b
Lymphocytes %	4.72 \pm 0.75 ac	4.22 \pm 0.54 a	6.06 \pm 1.44 b	5.52 \pm 0.62 bc
Monocytes %	1.84 \pm 0.52 a	1.46 \pm 0.34 a	1.8 \pm 0.32 a	1.56 \pm 0.30 a
Granulocytes %	1.17 \pm 0.14 a	1.38 \pm 0.36 ac	1.6 \pm 0.33 ac	1.7 \pm 0.23 bc
Platelets (cell $\times 10^9\mu\text{L}$)	139.6 \pm 8.32 a	126.2 \pm 12.54 b	219.8 \pm 27.62 c	152 \pm 12.63 a

In this study, increase of hemoglobin in HG and HSG can be attributable to the high concentration of minerals, particularly iron, and ascorbic acid in *H. sabdariffa* (Da-Costa et al., 2014). Previous studies showed that *H. sabdariffa* improve iron status and increase hemoglobin in women (Kubuga et al., 2019).

Lymphocytes and granulocytes increased in *H. sabdariffa* groups in our study. The presence of phytochemicals like flavonoids and tannins in the *H. sabdariffa* extract could have hemopoietic stimulating effect (Al-Jarah et al., 2017). Studies identified effect of *H. sabdariffa* in improve of hematological parameters such as RBC, WBC, Hb, and PCV in Swiss albino mice (Bhakta and Das, 2019). This evidence could boost immune system and defensive mechanism in the body.

In this study, platelets increased in *H. sabdariffa* group, which could be attributable to presence of components that produce blood cells such as vitamin C, thiamin and riboflavin, as well as β -carotene that transform to vitamin A, a helpful vitamin in treatment of platelet disorder (Carvajal-Zarrabal et al., 2012). In our study, on the other hand, platelets decreased in rabbits treated with starvation. Feng et al (2019) showed that autophagosome formation induced in platelets by starvation because of unwanted activation of platelets during starvation.

Liver Function


Liver function test in this study revealed that ALT activity significantly increased in SG, HG, and HSG groups, compared to CG group, although it was the same in both HG and HSG groups (Table 2). Moreover, AST activity significantly increased in both HG and HSG groups, compared to CG and SG groups (Table 2). In addition, histopathological sections of the liver revealed changes in the liver tissue in SG, HG, and HSG (Fig. 1, 2, 3). The Histopathological changes included coagulative necrosis of hepatocytes, hyperplasia of bile duct endothelia, dilation and congestion of central veins, vacuolar degeneration of the

hepatocytes around the central veins, and presence of inflammatory cells, particularly macrophages, around the central veins.


Increase the activity of ALT and AST in the circulation indicates hepatocellular damage, toxicity, inflammation, hypoxia, or tissue trauma (Igwilo et al., 2019; Huang et al., 2015; Chapman and Hostutler, 2013). It is known that inflamed or injured liver cells leak high amount of certain chemicals including ALT and AST into blood stream (Ejere et al., 2013; Esposti et al., 2012). A previous study showed that prolonged use of aqueous extract of calyx *H. sabdariffa* caused liver injury even at dose level lower than 150 mg/Kg B.W. (Ojokoh, 2010). In addition, ALT increased in starvation group, suggesting that alanine is an important amino acid involved in neoglucogenesis during starvation (Sharma et al., 1985). Changes in ALT and AST can suggest a rapid mobilization between intra- and extra hepatic fat stores that can cause histopathological changes (Gasteyger et al., 2017). Fasting can cause increase of hepatic content of triacylglycerol resulting in accumulation of lipid in the liver parenchyma that appear as vacuoles (Namazi et al., 2016; Lonardo et al, 2015; Marks et al., 2015).

Table 2. Differences in ALT and AST activity between study groups, control (CG); starvation (SG); Hibiscus *sabdariffa* extract (HG); and Hibiscus *sabdariffa* extract and starvation (HSG) groups, at the end of the study. Different letters (i.e., a, b, c, d) for each parameter refer to statistical difference ($P \leq 0.05$).


Liver function Parameter	Mean \pm Standard deviation			
	CG	SG	HG	HSG
ALT (U/L)	113.06 \pm 4.68 a	121.98 \pm 3.24 B	149.26 \pm 19.13 c	143.36 \pm 15.27 c
AST (U/L)	80.46 \pm 6.55 a	83.6 \pm 4.59 A	116.88 \pm 5.93 b	116.52 \pm 3.89 b



(Fig. 1) Histopathological section of liver from starvation group. Coagulative necrosis (a) of hepatocytes including pycnosis, karyorehexis, and karyolysis. There is hyperplasia of bile duct endothelia, dilation and congestion of central veins (b), and presence of inflammatory cells (c), particularly macrophages, around the central veins.



(Fig. 2) Histopathological section of liver from hibiscus extract group. Histopathological changes included dilation of central veins (a), vacuolar degeneration of the hepatocytes around the central veins (b), infiltration of inflammatory cells in the portal area (c), portal vein and hepatic artery congestion (d), and focal infiltration of inflammatory cells in the hepatic tissue and around central veins.




(Fig. 3) Histopathological section of liver from hibiscus extract with starvation group. Histopathological changes included vacuolar degeneration (a), coagulative necrosis, dilation of sinusoids, dilation of central veins (b) and infiltration of inflammatory cells in the portal area (c).

Lipid Profile

In this study, cholesterol and triglycerides decreased in SG, HG, and HSG groups, compared to CG group; however, there was no difference between SG and ESG groups (Table 3). In addition, no histopathological changes revealed in rabbits treated with *H. sabdariffa*. On the other hand, aorta sections from rabbits in starvation group revealed simple changes in aorta layers including few number of foam cells in tunica media, which appeared colorless as vacuoles.

Table 3. Differences in cholesterol and triglycerides concentrations between study groups, control (CG); starvation (SG); Hibiscus *sabdariffa* extract (HG); and Hibiscus *sabdariffa* extract and starvation (HSG) groups, at the end of the study. Different letters (i.e., a, b, c, d) for each parameter refer to statistical difference ($P \leq 0.01$).

Lipid profile parameter	Mean \pm Standard deviation			
	CG	SG	HG	HSG
Cholesterol (mg/dL)	134.1 \pm 9.34 a	115.04 \pm 6.17 B	99.63 \pm 12.85 c	119.46 \pm 1.99 B
Triglycerides (mg/dL)	153.22 \pm 3.54 a	137.77 \pm 4.43 B	121.16 \pm 4.05 c	121.80 \pm 3.76 C



(Fig. 4) Histopathological section of aorta from starvation group. Simple changes in aorta layers including few number foam cells (→) in tunica media, which appeared colorless as vacuoles.

In this study, the decrease in serum levels of cholesterol and triglycerides in rabbits treated with *H. sabdariffa* could be mediated by the antioxidant characteristics in the extract (Aba et al., 2016). Moreover, the *H. sabdariffa* extract is able to synthesis of HDL from the liver helping clearing excess cholesterol from the body in a process of inverse cholesterol transport (Elijah et al., 2017). Our result is similar to those reported by Aguirre-García et al (2019). In addition, Lin et al (2007) observed that *H. sabdariffa* can decrease serum cholesterol levels in human.

The normal histopathological sections in rabbits treated with *H. sabdariffa* extract could be attributable to the antioxidant effect of the extract that can protect the aorta from the oxidative damage and prevent development of atherosclerotic lesions (Zheoat et al., 2019; Zainalabidin et al., 2018; Yusof et al., 2018). On the other hand, a mild histopathological changes observed in starvation group that can be attributable to intracellular lipid mobilization as a result of utilizing the fat as a source of energy (Kartin et al., 1944). In this situation, a gradual accumulation of lipid into aortic subendothelial space occurs, which is taken by macrophage and transform into foam cells (Maguire et al., 2019). These changes are most likely prevented in rabbits with starvation and extract group as a result of antioxidant effect of the extract, an action that requires more investigation.

CONCLUSIONS

This study concluded that the aqueous *Hibiscus sabdariffa* can decrease the body weight as alternative to the starvation, decrease blood cholesterol and triglycerides, and increase the immunity; however, daily consumption can have impact on the liver.

REFERENCES

- Aba PE, Ossai A, Nweze EC. Preliminary Investigation of the Subchronic Hypolipidemic Actions of *Hibiscus sabdariffa* in Rats. Journal of Advances in Medical and Pharmaceutical Sciences, 2016;7(2):2394-1111. <https://doi.org/10.9734/JAMPS/2016/23186>
- Abbas MK, Ali AS. Effect of foliar application of NPK on some growth characters of two cultivars of Roselle (*Hibiscus sabdariffa* L.). American Journal of Plant Physiology; 2011;6(4),222-227.
- Aguirre-García F, Yáñez-López L, Armella MA, Verde JR. Studies from *Hibiscus sabdariffa* (Hibiscus) Plant for Blood Cholesterol Levels Reduction. American Journal of Plant Sciences, 2019;10:497-511 <https://doi.org/10.4236/ajps.2019.104036>
- Al-Jarah G, Mohammed A, and Jasim HA. Effects of Aqueous and Methanolic Extracts of *Hibiscus rosa-sinensis* Leave on the Blood Cells of Mice In vivo. World J Exp Biosci. 2017;5(1):46-49.
- Alka Sawarkar, Jangde CR, Thakre PD, Ranu K, and Shelu S. Analgesic Activity of *Hibiscus rosa sinensis* Linn in Rat. Veterinary World, 2009;2(9):353-354.
- Al-Snafi A. Pharmacological and therapeutic importance of *hibiscus sabdariffa*-A review. International Journal of Pharmaceutical Research. 2018;10:451-475.
- Bako IG, Mabrouk MA, Maje IM, Buraimoh AA and Abubakar MS. Hypotensive Effect of Aqueous Seed Extract of *Hibiscus sabdariffa* linn (*Malvaceae*) on Normotensive Cat. International Journal of Animal and Veterinary Advances, 2010;2(1):5-8.
- Bhakta S, and Das Sk. Effects of *Hibiscus Rosasinensis* on the Hematology of the Swiss Albino Mice: An In-Vivo. J App Mat Sci Engg Res, 2019;3(2)1 of 4.
- Carvajal-Zarrabal O, Barradas-Dermitz DM, Orta-Flores Z, et al. *Hibiscus sabdariffa* L., roselle calyx, from ethnobotany to pharmacology. J Exp Pharmacol. 2012;4:25–39. <https://doi.org/10.2147/JEP.S27974>
- Chapman SE, Hostutler RA. A Laboratory Diagnostic Approach to Hepatobiliary Disease in Small Animals. Veterinary Clinics of North America: Small Animal Practice. 2013;43(6),1209-1225. <https://doi.org/10.1016/j.cvsm.2013.07.005>
- Chen, C, Chou F, Ho Y, Lin W, Wang C, Kao E, Huang A, and Wang Ch. Inhibitory effects of *Hibiscus sabdariffa* L extract on low-density lipoprotein oxidation and anti-hyperlipidemia in fructose-fed and cholesterol-fed rats. Journal of the Science of Food and Agriculture. 2004;84:1989-1996. <https://doi.org/10.1002/jsfa.1872>.
- Da-Costa-Rocha I, Bonnlaender B, Sievers H, Pischel I, Heinrich M. *Hibiscus sabdariffa* L. – A phytochemical and pharmacological review. Food Chem., 2014;165:424-43. <https://doi.org/10.1016/j.foodchem.2014.05.002>.
- Drury RAB, Wallgton EA, and Cameron SR. Carleton's histological techniques. 4th ed. Oxford Univ. Press. New York. 1985.

- Duke J. and Atchley A. Proximate Analysis. In: The Hand Book of Plant Science in Agricultural. Christie, B.R. (ed.). CRC Press, Inc., Boca Raton Florida, USA; 1984.
- Ejere VC, Nnamonu DI, Chukwuka CO, Ugwu GD, Ejim AO, Asogwa CN. Effects of aqueous extract of *Hibiscus sabdariffa* calyces on haematological characteristics of *Rattus novergicus*. *Animal Research International*, 2013;10(3):1809-1816.
- Elijah JP, Young UC, Chuma C, Chinelo NC, Onyiaoziri OJ, Ogara A, Obiageli NF, Vincent O, Oteri EJ, and Emmanuela UN. Biochemical Effects of the Aqueous Extract of *Hibiscus sabdariffa* on Liver Marker Enzymes and Lipid Profiles in Acetaminophen-challenged Rats. *Research Journal of Medicinal Plants*, 2017;11:48-54. <https://doi.org/10.3923/rjmp.2017.48.54>
- Esposti DD, Hamelin J, Bosselut N, Saffroy R, Sebahg M, Pommier A, Martel C, Lemoine A. Mitochondrial roles and cytoprotection in chronic liver injury. *Biochem Res Int*. 2012;2012:387626. <https://doi.org/10.1155/2012/387626>
- Feng W, Chang C, Luo D, et al. Dissection of autophagy in human platelets. *Autophagy*. 2014;10(4):642–651. <https://doi.org/10.4161/auto.27832>
- Gasteyger C, Larsen TM, Vercruysse F, Astrup A. Effect of a dietary-induced weight loss on liver enzymes in obese subjects. *Am J Clin Nutr*. 2008;87(5):1141-7 <https://doi.org/10.1093/ajcn/87.5.1141>
- Gibbon D and Pain A. *Crops of the Drier Regions of the Tropics*. 1st ed. Longman Publishing Company, Singapore. 1985;155.
- Herranz-López, M., Olivares-Vicente, M., Boix-Castejón, M. et al. Differential effects of a combination of *Hibiscus sabdariffa* and *Lippia citriodora* polyphenols in overweight/obese subjects: A randomized controlled trial. *Sci Rep*. 2019; <https://doi.org/10.1038/s41598-019-39159-5>
- Hirunpanich V., Utaipat A., Morales N.P., Bunyaphatsara N., Sato H., Herunsalee A., Suthisang C. (2005). Antioxidant effects of aqueous extracts from dried calyx of *Hibiscus sabdariffa* Linn. (Roselle) in vitro using rat low-density lipoprotein (LDL). *Biol Pharm Bull*. 28(3):481-4 <https://doi.org/10.1248/bpb.28.481>
- Huang T-W, Chang C-L, Kao E-S, Lin J-H. Effect of *Hibiscus sabdariffa* extract on high fat diet-induced obesity and liver damage in hamsters. *Food Nutr Res*. 2015;15:59:29018. <https://doi.org/10.3402/fnr.v59.29018>
- Hussein RM, Yasser E, Shahein YE, Amr E, El Hakim AE, Hanem M, et al. Biochemical and molecular characterization of three colored types of roselle (*Hibiscus sabdariffa* L.) J. *Am. Sci*. 2010;11:726–733.
- Igwilo PC, Aguzie ION, Agbawodike CR, Attamah GN, Okafor FC. Hepatoprotective Activities of Methanol Extract of *Hibiscus sabdariffa* Lin. in Paracetamol-induced Hepatotoxicity of Albino Rats. *IOSR-JPBS*, 2019;14(4):01-08.

- Ismail A, Ikram EHK, Nazri HSM. Roselle (*Hibiscus sabdariffa* L.) seeds nutritional composition protein quality and health benefits. *Food*, 2008;2(1):1-16.
- Jiménez-Ferrer E1, Alarcón-Alonso J, Aguilar-Rojas A, Zamilpa A, Jiménez-Ferrer CI, Tortoriello J, Herrera-Ruiz M. Diuretic effect of compounds from *Hibiscus sabdariffa* by modulation of the aldosterone activity. *Planta Med.* 2012;78(18):1893-8. <https://doi.org/10.1055/s-0032-1327864>
- Kartin BL, Man EB, Winkler AW, Peters JP. Blood Ketones And Serum Lipids In Starvation And Water Deprivation. *J Clin Invest.* 1944;23(5):824–835. <https://doi.org/10.1172/JCI101556>
- Kaur C, and Kapoor HC. Antioxidant activity of some fruits in Indian diet. *Acta Hort.*, 2005; 563-565. <https://doi.org/10.17660/ActaHortic.2005.696.99>
- Kochhar SL Okra (Lady's finger) In: *Tropical crops, a textbook of economic Botany*. Editor S.L., Kochhar, 1986;263-264.
- Kubuga K, Hong HG, and Song WO. *Hibiscus sabdariffa* meal improves iron status of childbearing age women and prevents stunting in their toddlers in northern Ghana. *Nutrients* 2019;11:198 <https://doi.org/10.3390/nu11010198>
- Lin TL, Lin HH, Chen CC, Lin MC, Chou MC, Wang CJ. *Hibiscus sabdariffa* extract reduces serum cholesterol in men and women. *Nutr Res*, 2007;27(3): 140-145 <https://doi.org/10.1016/j.nutres.2007.01.007>
- Lonardo A, Ballestri S, Marchesini G. Nonalcoholic fatty liver disease: A precursor of the metabolic syndrome. *Digest. Liver Dis.* 2015;47(3): 181-190.
- Luna LG. *Manual of histologic staining methods of the Armed Forces Institute of Pathology*. 3rd ed, McGraw-Hill, New York. 1968.
- Madea B. *Death as a Result of Starvation Part of the Forensic Pathology Reviews book series (FPR, volume 2)*. Edited by: M. Tsokos © Humana Press Inc., Totowa, NJ, 2005.
- Maguire EM, Pearce SWA, Xiao Q. Foam cell formation: A new target for fighting atherosclerosis and cardiovascular disease. *Vascul Pharmacol.*, 2019;112:54-71. <https://doi.org/10.1016/j.vph.2018.08.002>
- Mahadevan NS, Kamboj P. *Hibiscus sabdariffa* Linn.-An overview. *Natural Product Radiance*, 2009;8(1):77-8.
- Marks KA, Marvyn PM, Henao JJ, Bradley RM, Stark KD, Duncan RE. Fasting enriches liver triacylglycerol with n-3 polyunsaturated fatty acids: implications for understanding the adipose-liver axis in serum docosahexaenoic acid regulation. *Genes Nutr*, 2015;10(6):1-14. <https://doi.org/10.1007/s12263-015-0490-2>
- McKay DL, Chen CY, Saltzman E, Blumberg JB. *Hibiscus Sabdariffa* L. Tea (Tisane) Lowers Blood Pressure in Prehypertensive and Mildly Hypertensive Adults. *J Nutr.* 2010;140(2):298-303. <https://doi.org/10.3945/jn.109.115097>

- Moore, D. S., Craig, B. A., & McCabe, G. P. (2009). Introduction to the practice of statistics. 6th ed. New York, NY: W. H. Freeman and Co. Pp: 450-455.
- Mungole A, and Chaturvedi A. *Hibiscus sabdariffa* L a rich source of secondary metabolites. International Journal of Pharmaceutical Sciences Review and Research. 2011;6:83-87.
- Namazi F, Omidi A, Abbasi S, Afsar M, Honarmand M, and Nazifi S. Starvation and refeeding in rats: effect on some parameters of energy metabolism and electrolytes and changes of hepatic tissue. Pesq Vet Bras 2016;36(1):101-105 <http://dx.doi.org/10.1590/S0100-736X2016001300015>
- Odigie IP, Ettarh RR, and Adigun SA. Chronic administration of aqueous extract of *Hibiscus Sabdariffa* attenuates hypertension and reverses cardiac hypertrophy in 2K-1C hypertensive rats. J Ethnopharmacol, 2003;86(2-3):181-5.
- Ojeda D, Jiménez-Ferrer E, Zamilpa A, Herrera-Arellano A, Tortoriello J, Alvarez L. Inhibition of angiotensin convertin enzyme (ACE) activity by the anthocyanins delphinidin- and cyanidin-3-O-sambubiosides from *Hibiscus sabdariffa*. Nutr Metab (Lond). 2009;2(6):45. <https://doi.org/10.1186/1743-7075-6-45>
- Ojokoh AO. Roselle (*Hibiscus Sabdariffa*) Calyx Diet and Histopathological Changes in Liver of Albino Rats. Pakistan Journal of Nutrition, 2006;5(2)110-113 <https://doi.org/10.3923/pjn.2006.110.113>
- Sanchez-Elsner T, Ramirez JR, Rodriguez-Sanz F, Varela E, Bernabew C, Botella LM. A cross talk between hypoxia and TGF-beta orchestrates erythropoietin gene regulation through SPI and smads. J Mol Biol. 2004;36(1):9–24.
- Sarr M, Ngom S, Kane MO, Wele A, Diop D, Sarr B, Gueye L, Andriantsitohaina R, Diallo AS. In vitro vasorelaxation mechanisms of bioactive compounds extracted from *Hibiscus sabdariffa* on rat thoracic aorta. 2009;2(6):45. <https://doi.org/10.1186/1743-7075-6-45>
- Sharma M, Lal H, Saini AS. Effect of prolonged starvation and refeeding on fuel metabolism in rats. Indian J Physiol Pharmacol. 1985;29(2):107-10.
- Singh P, Khan M, Hailemariam H. Nutritional and Health Importance of *Hibiscus Sabdariffa*: A Review and Indication for Research Needs. J Nutr Health Food Eng; 2017;6(5): 00212. <https://doi.org/10.15406/jnhfe.2017.06.00212>
- Voon TB, Mole P, Banca L, Porter L, Morris S, Mitchell S, Lapa TR, Karr J, Harrison NA, Potenza MN, Irvine M. Neural correlates of sexual cue reactivity in individuals with and without compulsive sexual behaviours. PLoS One. 2014;9(7):e102419. <https://doi.org/10.1371/journal.pone.0102419>
- Yusof NLM, Jubaidi FF, Nasir SNM, Yusoff NA, Mohd Fauzi N, Zainalabidin S, and Budin SB. *Hibiscus sabdariffa* (Roselle) polyphenol-rich extract prevents the aortic oxidative damage in type 1 diabetic rats. Jurnal Teknologi, 2018;80(2):1-8. <https://doi.org/10.11113/jt.v80.11165>

Zahid H, Rizwani GH, Shareef H, Khursheed R, Huma A, Hasan SM. Antihyperglycemic and hypolipidemic effects of *Hibiscus schizopetalus* (Mast) Hook in alloxan-induced diabetic rats. *Pak J Pharm Sci.* 2014;27(1):83-9.

Zainalabidin S, Budin SB, Mohamad Anuar NN, Yusoff NA, and Yusof NLM. *Hibiscus sabdariffa* Linn. improves the aortic damage in diabetic rats by acting as antioxidant. *Journal of Applied Pharmaceutical Science*, 2018;8(1):108-114. <https://doi.org/10.7324/JAPS.2018.8116>

Zheoat AM, Gray AI, Igoli JO, Ferro VA, Drummond RM. Hibiscus acid from *Hibiscus sabdariffa* (Malvaceae) has a vasorelaxant effect on the rat aorta. *Fitoterapia*, 2019;134:5-13. <https://doi.org/10.1016/j.fitote.2019.01.012>